

## **A Molecular Dialogue Between the Brain and Face**

Ralph Marcucio, Dwight Cordero, and Diane Hu, and **Jill A. Helms\*** University of California at San Francisco, CA USA 94143-0514. Telephone: 415 502 6523; Fax: 415 476 1128; Email: helms@cgl.ucsf.edu

Clinicians have long recognized that a relationship exists between the development of the forebrain and the face. Forebrain defects are frequently associated with distinctive facial features [1], allowing experienced clinicians to diagnose brain anomalies based on a patient's facial appearance. Traditionally the forebrain has been viewed as a scaffold upon which the face develops, and this observation has led to the notion that forebrain defects are accompanied by facial defects due primarily to mechanical influences of one tissue on the other [2].

While the brain plays a substantial physical role in shaping the face, the forebrain and face are also intimately linked because morphogenesis of both structures requires tissue interactions among forebrain neuroepithelium, neural crest, and ectoderm of the face [3]. All of the facial primordia are populated by mesenchymal cells derived principally from the neural crest and are surrounded by epithelia of ectodermal and in some cases, endodermal origin. Through a series of poorly understood molecular and cellular events, these undifferentiated buds of tissue are transformed into the anatomically intricate forehead, nose, cheeks, lips, jaws, and chin. How do cells within the developing facial primordia acquire the spatial information necessary to produce such highly integrated morphological components? We propose that regions of neural and facial epithelia act as organizing centers, which mediate dorsoventral, mediolateral, and proximodistal patterning of the FNP through the regulation of bone morphogenetic proteins (Bmps). We will provide data demonstrating that frontonasal process (FNP) ectodermal cues specify dorsoventral patterning in the mid- and upper face. We will also show data indicating that the neuroepithelial domain of Sonic hedgehog (Shh; 4) establishes the mediolateral axis of the FNP, in part by mediating migration of FNP neural crest to the facial midline. Collectively, these data offer insight into the mechanisms by which epithelial-mesenchymal interactions govern development of the mid- and upper-face, and provide a fundamental framework upon which preventive, diagnostic, and therapeutic approaches to treating birth defects can be developed.

### **References:**

1. Gorlin, R.J., M.M. Cohen, and L.S. Levin, Syndromes of the Head and Neck. 3rd ed. Oxford Monographs on Medical Genetics. Vol. 1. 1990, New York: Oxford University Press.
2. DeMyer, W., The face predicts the brain: diagnostic significance of median facial anomalies for holoprosencephaly (arhinencephaly). *Pediatrics*, 1964. August: p. 256-263.
3. Schneider, R.A., et al., Local retinoid signaling coordinates forebrain and facial morphogenesis by maintaining FGF8 and SHH. *Development*, 2001. 128(14).
4. Riddle, R.D., et al., Sonic hedgehog mediates the polarizing activity of the ZPA. *Cell*, 1993. 75(7): p. 1401-16.

### **Reviews:**

R. Schneider, D. Hu and J. A. Helms. From head to toe: the conservation of molecular signals regulating limb and craniofacial morphogenesis. *Cell and Tissue Research* 296:103-9, 1999. K. Uniscker, editor.

D. Young, R. Schneider and J. A. Helms. Genetic and teratogenic approaches to craniofacial development. Annual reviews in Oral Medicine and Biology, 2000;11(3):304-17.

**Original articles:**

D. Hu and J. A. Helms. Patterning the face: the role of Sonic hedgehog in normal and abnormal craniofacial morphogenesis. Development 126:4873-4884, 1999.

K. Suzuki, D. Hu, T. Bustos, J. Zlotogora, A. Richieri-Costa, J.A. Helms, R.A. Spritz. Mutations of PVRL1 cell-cell adhesion molecule/Herpesvirus receptor gene in cleft lip/palate-ectodermal dysplasia. Nature Genetics 25:427-30, 2000.

R.A. Schneider, D. Hu, J. Rubenstein, M. Maden, and J.A. Helms. Local retinoid signaling coordinates forebrain and facial morphogenesis by maintaining FGF8 and SHH. Development, 128(14), 2001.